

Maps of the **published** cases of resistance to herbicides in France

Maps of the **published** cases of resistance to herbicides in France

Foreword

1. The herbicide classification used in this version is the one developed by R4P for **all** pesticides

<https://www.r4p-inra.fr/fr/classification-des-ppp/>

2. The weed species mapped are exclusively those for which there is at least one published paper describing the resistance case, or for which resistance has been demonstrated in the national network of Biological Surveillance of the Territory (SBT).

➔ **the list is not exhaustive**, as some existing cases remain to be published.

3. On the maps, one French department is coloured according to the number of resistance cases reported.

➔ Obviously, this does not mean that all the fields in a coloured department are affected by resistance, even for departments where resistance is very frequent!

➔ Conversely, a department not being coloured only means that resistance has not been reported (either because it is really absent, or because it has not been searched or identified, or because the data have not been communicated).

Resistance case summary: the family picture

Legend to the pictures:

Weed

Herbicide mode(s) of action concerned by resistance (R4P classification) (year when resistance has been first reported)

Crops infested by the weed

Prevalence of the resistance in the crops concerned (first reports, moderate, common, very common)

Grass weeds I



Black-grass

D1 (1993),
F2 (2006)
Very common
D2 (2019)
F1 (2022)
First report
Major crops



Bromes

F2 (2009)
Major crops
Moderate



Rye-grasses

D1 (1993),
F2 (2001),
Very common
D2 (2014)
First reports
Major crops
F1 (2005)
Grapevine, major crops
Moderate



Wild oats

D1 (1996),
F2 (2006 approx.)
Major crops
Moderate



Silky bent-grass

D1 (1999 approx.),
F2 (2006 approx.)
Major crops
Moderate



Large crabgrass

D1 (2005)
Vegetables
F2 (2014)
I2 (2020)
Maize
First reports



Foxtails

F2 (2011) Maize
First reports



Barnyard-grass

D1 (2000 approx.)
Rice
Very common
F2 (2013)
Rice
Very common
Maize
First reports
I2 (2017)
Maize
First reports



Rice barnyard-grass

D1 (2000 approx.)
F2 (2013)
Rice
Very common



Rough meadow-
grass
F2 (2012)
Major crops
First reports



Annual
bluegrass
D1 (not
sensitive)
F2 (2015)
Major crops
First reports




Teosinte
D1 (1995s)
Maize, soybean,
sunflower
Moderate



Common groundsel
F2 (2009)
Grapevine, major crops, vegetables...
Common



Weedy sunflower
F2 (2009)
ALS-tolerant sunflower, soybean
Moderate



Chamomiles
F2 (2010)
Major crops
Moderate



Sumatra fleabane
F1 (2010)
F2 (2016)
Grapevine
Moderate



Common ragweed
F2 (2014)
Soybean, ALS-tolerant sunflower, maize
Moderate



Spiny sow-thistle
F2 (2016)
Chicory
Very common



Gallant soldier
F2 (2017)
Chicory
First reports



Canadian horseweed
F1 (2019)
Grapevine
Moderate



Corn poppy
F2 (2007)
Major crops
Very common
L1 (2013)
Major crops
Moderate



Common lambsquarters
B1 (2005)
Sugar beet,
potatoe
Common



Chickweed
F2 (2012)
Major crops
First reports



Bitter dock
F2 (2018)
Major crops
First reports



Spear saltbush
B1 (2019)
Sugar beet,
potatoe
First reports

To sum it all:

Resistance:

- To graminicide herbicides – ACCase inhibitors (group D1)
 - To ALS inhibitors (group F2)
 - To « auxinic » herbicides (group L1)
- To some root-applied herbicides in group D2
 - To photosystem II inhibitors (group B1)
 - To glyphosate (EPSPS inhibitor, group F1)
 - To HPPD inhibitors (group I2)

Grass weeds & Asteraceae

(+ poppy, lambsquarter, speat saltbush, bitter dock & chickweed)

Updated regularly on:

www.r4p-inra.fr

Maps

Resistance to herbicides in France: **published** cases

Grass weeds


Resistance of black-grass to ACCase inhibitors (group D1)

Updated: October 2026



Alopecurus myosuroides

- Common (> 20 cases)
- Moderate (5-20 cases)
- Rare (2-4 cases)
- First case reported
- Not reported



Resistance: **NTSR** + **TSR**

Cereals, rape, sugar beet...

Actives concerned: fenoxaprop, clodinafop, fluazifop, propaquizafop, quizalofop, pinoxaden, cycloxydim, clethodim...

Data: INRAE, COLUMA

Multiple resistance / cross-**NTSR** with herbicides in groups D2 and/or F2 and/or F1


Resistance of black-grass to ALS inhibitors (group F2)

Updated: October 2026



Alopecurus myosuroides

- Common (> 20 cases)
- Moderate (5-20 cases)
- Rare (2-4 cases)
- First case reported
- Not reported



Resistance: **NTSR** + **TSR**

Cereals, rape...

Actives concerned: iodosulfuron + mesosulfuron, pyroxsulam, imazamox...

Data: INRAE, COLUMA

Multiple resistance / cross-**NTSR** with herbicides in groups D1 and/or D2 and/or F1


Resistance of black-grass to inhibitors of lipid elongation (group D2)

Updated: October 2026



Alopecurus myosuroides

- Common (> 20 cases)
- Moderate (5-20 cases)
- Rare (2-4 cases)
- First case reported
- Not reported



Resistance: mechanism not investigated (probably **NTSR**)

Cereals...

Actives concerned: flufenacet, prosulfocarb

Data: INRAE

Multiple resistance / cross-**NTSR** with herbicides in groups D1 and/or F2 and/or F1


Resistance of black-grass to inhibitors EPSPS (group F1)

Updated: October 2026



Alopecurus myosuroides

- Common (> 20 cases)
- Moderate (5-20 cases)
- Rare (2-4 cases)
- First case reported
- Not reported



Resistance: mechanism not investigated

Cereals

Active concerned: glyphosate

Data: INRAE, SBT

Multiple resistance with herbicides in groups D1 and/or D2 and/or F2


Resistance of rye-grasses to ACCase inhibitors (group D1)

Updated: October 2026



Lolium sp.

- Common (> 20 cases)
- Moderate (5-20 cases)
- Rare (2-4 cases)
- First case reported
- Not reported



Resistance: **NTSR** + **TSR**

Cereals, rape, sugar beet...

Actives concerned: clodinafop, fluazifop, propaquizafop, quizalofop, pinoxaden, cycloxydim, clethodim...

Data: INRAE, COLUMA

Multiple resistance / cross-**NTSR** with herbicides in groups D2 and/or F2 and/or F1


Resistance of rye-grasses to ALS inhibitors (group F2)

Updated: October 2026



Lolium sp.

- Common (> 20 cases)
- Moderate (5-20 cases)
- Rare (2-4 cases)
- First case reported
- Not reported



Resistance: **NTSR** + **TSR**

Cereals, rape...

Actives concerned: iodosulfuron + mesosulfuron, pyroxsulam, imazamox...

Data: INRAE, COLUMA

Multiple resistance / cross-**NTSR** with herbicides in groups D1 and/or D2 and/or F1


Resistance of rye-grass to inhibitors of lipid elongation (group D2)

Updated: October 2026



Lolium sp.

- Common (> 20 cases)
- Moderate (5-20 cases)
- Rare (2-4 cases)
- First case reported
- Not reported



Resistance: mechanism not investigated (probably **NTSR**)

Cereals...

Actives concerned: flufenacet, prosulfocarb

Data: INRAE, Bayer, SBT

Multiple resistance / cross-**NTSR** with herbicides in groups D1 and/or F2 and/or F1


Resistance of rye-grasses to EPSPS inhibitors (group F1)

Updated: October 2026



Lolium sp.

- Common (> 20 cases)
- Moderate (5-20 cases)
- Rare (2-4 cases)
- First case reported
- Not reported



Resistance: mechanism not investigated / altered translocation (**NTSR**) + **TSR**

Cereals and grapevine

Active concerned: glyphosate

Note: the data for grapevine dates back to 2012...

Multiple resistance with herbicides in groups D1 and/or D2 and/or F2

Data: INRAE, SBT, Bayer


Resistance of Bromes to ALS inhibitors (group F2)

Updated: October 2026



Bromus sterilis

- Common (> 20 cases)
- Moderate (5-20 cases)
- Rare (2-4 cases)
- First case reported
- Not reported



Resistance: **TSR** + **NTSR**

Cereals, rape

Actives concerned: sulfosulfuron,
iodosulfuron + mesosulfuron,
pyroxsulam, imazamox,
propoxycarbazone...

Data: COLUMA


Resistance of wild oats to ACCase inhibitors (group D1)

Updated: October 2026



Avena sp.

- Common (> 20 cases)
- Moderate (5-20 cases)
- Rare (2-4 cases)
- First case reported
- Not reported



Resistance: **TSR + NTSR**

Cereals, rape

Actives concerned: fenoxaprop, clodinafop, fluazifop, propaquizafop, quizalofop, pinoxaden, cycloxydim, clethodime...

Data: COLUMA


Resistance of wild oats to ALS inhibitors (group F2)

Updated: October 2026



Avena sp.

- Common (> 20 cases)
- Moderate (5-20 cases)
- Rare (2-4 cases)
- First case reported
- Not reported



Resistance: **TSR** + **NTSR**

Cereals, rape

Actives concerned: iodosulfuron + mesosulfuron, pyroxsulam, imazamox...

Data: COLUMA


Resistance of silky bent-grass to ACCase inhibitors (group D1)

Updated: October 2026



Apera spica-venti

- Common (> 20 cases)
- Moderate (5-20 cases)
- Rare (2-4 cases)
- First case reported
- Not reported



Resistance: **TSR** + **NTSR**

Cereals, rape


Actives concerned: fenoxaprop, clodinafop, fluazifop, propaquizafop, quizalofop, pinoxaden, cycloxydim, clethodime...

Data: COLUMA



Apera spica-venti

- Common (> 20 cases)
- Moderate (5-20 cases)
- Rare (2-4 cases)
- First case reported
- Not reported



Resistance: **TSR** + **NTSR**

Cereals, rape

Actives concerned: iodosulfuron + mesosulfuron, pyroxsulam, imazamox...


Data: COLUMA

Resistance of barnyard-grass to ACCase inhibitors (group D1)



Echinochloa crus-galli

- Common (> 20 cases)
- Moderate (5-20 cases)
- Rare (2-4 cases)
- First case reported
- Not reported



Resistance: **TSR** (**NTSR?**)

Rice


Actives concerned: cycloxydim,
cyhalofop

Data: CFR



Echinochloa crus-galli

- Common (> 20 cases)
- Moderate (5-20 cases)
- Rare (2-4 cases)
- First case reported
- Not reported



Resistance: **TSR** (**NTSR?**)

Rice, maize

Actives concerned: azimsulfuron, prosulfuron, foramsulfuron, nicosulfuron, penoxsulam...

Resistance of barnyard-grass to HPPD inhibitors (group I2)


Updated: October 2026



Echinochloa crus-galli

- Common (> 20 cases)
- Moderate (5-20 cases)
- Rare (2-4 cases)
- First case reported
- Not reported

World premiere!



Resistance: mechanism not investigated
Maize
Actives concerned: mesotrione...


Data: INRAE

Resistance of rice barnyard-grass to ACCase inhibitors (group D1)



Echinochloa oryzicola

- Common (> 20 cases)
- Moderate (5-20 cases)
- Rare (2-4 cases)
- First case reported
- Not reported



Resistance: TSR (NTSR?)

Rice

Actives concerned: cycloxydim,
cyhalofop


Data: CFR

Resistance of rice barnyard-grass to ALS inhibitors (group F2)



Echinochloa oryzicola

- Common (> 20 cases)
- Moderate (5-20 cases)
- Rare (2-4 cases)
- First case reported
- Not reported



Resistance: **TSR** (**NTSR?**)

Rice

Actives concerned: azimsulfuron,
penoxsulam


Resistance of foxtails to ALS inhibitors (group F2)

Updated: October 2026



Setaria viridis

- Common (> 20 cases)
- Moderate (5-20 cases)
- Rare (2-4 cases)
- First case reported
- Not reported



Resistance: **TSR** (**NTSR?**)

Maize

Actives concerned: prosulfuron, foramsulfuron, nicosulfuron...


Data: INRAE, SBT

Resistance of large crabgrass to ACCase inhibitors (group D1)



Digitaria sanguinalis

- Common (> 20 cases)
- Moderate (5-20 cases)
- Rare (2-4 cases)
- First case reported
- Not reported



Resistance: **TSR**

Vegetables


Actives concerned: cycloxydim, clethodim, fluazifop, quizalofop...

Data: COLUMA



Digitaria sanguinalis

- Common (> 20 cases)
- Moderate (5-20 cases)
- Rare (2-4 cases)
- First case reported
- Not reported



Resistance: **NTSR** (**TSR?**)

Maize

Actives concerned: prosulfuron, foramsulfuron, nicosulfuron...

Resistance of large crabgrass to HPPD inhibitors (group I2)


Updated: October 2026



Digitaria sanguinalis

- Common (> 20 cases)
- Moderate (5-20 cases)
- Rare (2-4 cases)
- First case reported
- Not reported

World premiere!




Resistance: mechanism not investigated
Maize
Actives concerned: mesotrione...

Resistance of rough meadow-grass to ALS inhibitors (group F2)



Poa trivialis

- Common (> 20 cases)
- Moderate (5-20 cases)
- Rare (2-4 cases)
- First case reported
- Not reported



Resistance: **TSR**

Cereals


Actives concerned: iodosulfuron + mesosulfuron, pyroxsulam...

Data: Bayer



Poa annua

- Common (> 20 cases)
- Moderate (5-20 cases)
- Rare (2-4 cases)
- First case reported
- Not reported



Resistance: **TSR?**

Cereals, maize...

Actives concerned: iodosulfuron + mesosulfuron, pyroxsulam, prosulfuron, foramsulfuron, nicosulfuron ...

Note: this species is naturally little or not sensitive to ACCase inhibitors (group D1)

Data: Bayer


Resistance of Teosinte to ACCase inhibitors (group D1)

Updated: October 2026



Zea mays ssp. mexicana

- Common (> 20 cases)
- Moderate (5-20 cases)
- Rare (2-4 cases)
- First case reported
- Not reported



Resistance: TSR (by gene flow from herbicide-tolerant maize)

ACCase-tolerant maize, soybean, sunflower

Active concerned: cycloxydim

Data: INRAE

Broadleaves- Papaveraceae


Resistance of corn poppy to ALS inhibitors (group F2)

Updated: October 2026



Papaver rhoeas

- Common (> 20 cases)
- Moderate (5-20 cases)
- Rare (2-4 cases)
- First case reported
- Not reported



Resistance: **TSR** (+ **NTSR**)

Cereals, rape

Actives concerned: tribenuron, metsulfuron, tritosulfuron, iodosulfuron + mesosulfuron, florasulam, imazamox...


Multiple resistance with herbicides in group L1

Data: INRAE, Anses, COLUMA, SBT, Corteva



Papaver rhoeas

- Common (> 20 cases)
- Moderate (5-20 cases)
- Rare (2-4 cases)
- First case reported
- Not reported



Resistance: mechanism not investigated (NTSR?)

Cereals, rape

Actives concerned: 2,4-D, MCPA, aminopyralid...

Data: INRAE, Corteva,

Multiple resistance with herbicides in group F2


Broadleaves - Asteraceae

Resistance of common groundsel to ALS inhibitors (group F2)



Senecio vulgaris

- Common (> 20 cases)
- Moderate (5-20 cases)
- Rare (2-4 cases)
- First case reported
- Not reported



Resistance: **TSR** (+ **NTSR**)

Major crops, grapevine, chicory, vegetables


Actives concerned: tribenuron, metsulfuron, iodosulfuron + mesosulfuron, pyroxsulam, florasulam, imazamox, thiencarbazone...

Data: INRAE, Bayer, SBT, Corteva



Erigeron sumatrensis

- Common (> 20 cases)
- Moderate (5-20 cases)
- Rare (2-4 cases)
- First case reported
- Not reported



Resistance: **TSR** (**NTSR?**)

Grapevine

Actives concerned: flazasulfuron,
penoxsulam...

Multiple resistance with herbicides in group F1


Data: INRAE, IFVV,
Belchim, SBT

Resistance of Sumatra fleabane to EPSPS inhibitors (group F1)



Erigeron sumatrensis

- Common (> 20 cases)
- Moderate (5-20 cases)
- Rare (2-4 cases)
- First case reported
- Not reported



Resistance: mechanism not investigated

Grapevine

Active concerned: glyphosate

Data: INRAE, Monsanto, SBT

Multiple resistance with herbicides in group F2

Resistance of Canadian horseweed to EPSPS inhibitors (group F1)


INRAE

Updated: October 2026



Erigeron canadensis

- Common (> 20 cases)
- Moderate (5-20 cases)
- Rare (2-4 cases)
- First case reported
- Not reported



Resistance: mechanism not investigated

Grapevine


Active concerned: glyphosate

Data: INRAE, SBT



Helianthus annuus

- Common (> 20 cases)
- Moderate (5-20 cases)
- Rare (2-4 cases)
- First case reported
- Not reported



Resistance: TSR (by gene flow from herbicide-tolerant sunflower)


ALS-tolerant sunflower, soybean

Actives concerned: imazamox, tribenuron



Matricaria sp.

- Common (> 20 cases)
- Moderate (5-20 cases)
- Rare (2-4 cases)
- First case reported
- Not reported



Resistance: TSR (& NTSR?)

Cereals, rape

Actives concerned: tribenuron, metsulfuron, iodosulfuron + mesosulfuron, pyroxsulam, florasulam, imazamox...

Resistance of common ragweed to ALS inhibitors (group F2)

Updated: October 2026




-  Common (> 20 cases)
-  Moderate (5-20 cases)
-  Rare (2-4 cases)
-  First case reported
-  Not reported

Ambrosia artemisiifolia

Resistance: **NTSR** & **TSR**

ALS-tolerant sunflower, soybean, maize

Actives concerned: imazamox, tribenuron,...




Data: INRAE, BASF, SBT



Sonchus asper

- Common (> 20 cases)
- Moderate (5-20 cases)
- Rare (2-4 cases)
- First case reported
- Not reported



Resistance: TSR


Chicory (major crops)

Actives concerned: rimsulfuron,
triflusulfuron, penoxsulam, ...



Galinsoga parviflora

- Common (> 20 cases)
- Moderate (5-20 cases)
- Rare (2-4 cases)
- First case reported
- Not reported



Resistance: TSR

Chicory (major crops)

Actives concerned: rimsulfuron, triflusulfuron, penoxsulam, ...

Broadleaves - Others


Resistance of common lambsquarters to photosystem II inhibitors (group B1)

Updated: October 2026



Chenopodium album

- Common (> 20 cases)
- Moderate (5-20 cases)
- Rare (2-4 cases)
- First case reported
- Not reported



Resistance: **TSR**

Sugar beet, potatoe

Actives concerned:
metamitron, metribuzin...

Data: INRAE, ITB,
Corteva, Univ. Ghent


Resistance of spear saltbush to photosystem II inhibitors (group B1)

Updated: October 2026



Atriplex patula

- Common (> 20 cases)
- Moderate (5-20 cases)
- Rare (2-4 cases)
- First case reported
- Not reported



Resistance: TSR

Sugar beet, potatoe


**Actives concerned:
metamitron, metribuzin...**

Data: INRAE, ITB



Stellaria media

- Common (> 20 cases)
- Moderate (5-20 cases)
- Rare (2-4 cases)
- First case reported
- Not reported



Resistance: **TSR**

Cereals

Actives concerned: tribenuron, metsulfuron, iodosulfuron + mesosulfuron, pyroxsulam, florasulam, imazamox...


Resistance of bitter dock to ALS inhibitors (group F2)

Updated: October 2026



Rumex obtusifolius

- Common (> 20 cases)
- Moderate (5-20 cases)
- Rare (2-4 cases)
- First case reported
- Not reported



Resistance: **TSR**

Cereals

Actives concerned: metsulfuron, thifensulfuron, florasulam...

Data: FMC